

2018-5344
2019-05-17

GROUP	M	FUNGICIDE
-------	---	-----------

NET CONTENTS: 22.68 KILOGRAMS

BARTLETT MICROSCOPIC

Wettable Sulphur
AGRICULTURAL
ACTIVE INGREDIENT: SULPHUR 92%

REGISTRATION NO. 873 PEST CONTROL PRODUCTS ACT
READ THE LABEL BEFORE USING

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not inhale dust or spray mist. Avoid contact with skin and eyes. After handling, wash thoroughly with soap and water before eating or smoking. Do not contaminate ponds, streams or other bodies of water. Avoid contamination of feed or food. Wear goggles or a face shield, chemical-resistant gloves, a hat, a long-sleeved shirt, long pants and rubber boots during mixing/loading, application, clean-up and repair activities. Do not enter or allow workers to enter into treated areas until 24 hours after application. If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca.

FIRST AID: If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **In case of skin or clothing contact**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

DISPOSAL/DECONTAMINATION INSTRUCTIONS: 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. 2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal. 3. Make the empty container unsuitable for further use. 4. Dispose of the container in accordance with provincial requirements. 5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

RESISTANCE MANAGEMENT: For resistance management, please note that Microscopic Wettable Sulphur contains a Group M fungicide. Any fungal population may contain individuals naturally resistant to Sulphur and other Group M fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. To delay fungicide resistance:

- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for sign of resistance development.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact N.M. BARTLETT INC. at 1-800-263-1287 or at www.bartlett.ca.

COMPATIBILITY: This product is compatible with most commonly used insecticides and fungicides, except DNOC (Elgetol) and oil.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

DIRECTIONS FOR USE

The rates recommended below are based on 1000 L of dilute spray and kg per ha. Microscopic sulphur may be used in low volume or concentrate sprays by increasing the dosage to apply the same quantity of product per ha as would normally be applied when spraying dilute mixtures.

DO NOT APPLY BY AIR.

Follow provincial recommendations for proper timing of the sprays. **DO NOT APPLY IF HIGH TEMPERATURES (ABOVE 26°) AND HIGH HUMIDITY PREVAIL OR ARE EXPECTED DURING THE THREE DAYS FOLLOWING APPLICATION. DO NOT USE SULPHUR WITHIN 30 DAYS OF AN OIL SPRAY OR BORDEAUX MIXTURE.** Do not apply more than 8 applications per season. Do not apply during periods of dead calm or when winds are gusty. For airblast application: Do not direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. Do not apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

CROP/DISEASE	METHOD/RATE
APPLE, PEAR Powdery Mildew Scab	Apply 6.5 kg per 1000 L (19.5 kg per ha) in the Pre-Bloom, Calyx and first Cover sprays. Reduce to 3 kg/1000 L or 9.0 kg per ha if hot weather threatens. In the later Cover sprays use 3.5 kg - 4.5 kg (10.5 kg - 13.5 kg per ha) depending on mildew conditions and temperature.
PEACH Brown Rot Scab	Apply 6.5 kg per 1000 L (19.5 kg per ha) in a regular spray program starting at the late Pink stage. Repeat just before and during harvest if wet weather occurs.
PLUM, PRUNE Brown Rot Black Knot	Apply 6.5 kg per 1000 L (19.5 kg per ha) in a regular spray program starting at the Shuck stage. Repeat just before and during harvest if wet weather occurs.
SWEET CHERRY Brown Rot	Apply 6.5 kg per 1000 L (19.5 kg per ha) in a regular spray program from the Bloom stage until just before harvest. Repeat during harvest if wet weather occurs.
SOUR CHERRY Brown Rot Powdery Mildew	Apply 6.5 kg per 1000 L (19.5 kg per ha) at the bloom stage. If powdery mildew is a problem, apply 3.5 kg/1000 L (10.5 kg per ha) in the first, second and third cover spray.
GRAPES Powdery Mildew	Apply 4.5 kg/1000 L in the Post-Bloom spray on Agawam, Seneca and Seibel varieties. Do not use on Concord, Foch, Van Buren or other sulphur sensitive varieties of grapes. Apply 6 kg/1000 L in the First, Second and further Cover sprays. Do not apply later than 21 days before harvest on grapes for wine.

Table 1. Buffer zones for the protection of aquatic habitats. This table specifies the buffer zones which are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

METHOD OF APPLICATION	CROP	BUFFER ZONE (METRES) REQUIRED FOR THE PROTECTION OF AQUATIC HABITAT
Airblast	Sweet cherry Sour Cherry Plum Peach Apple Pear Prune Grape	5

TOXICOLOGICAL INFORMATION: Treat Symptomatically.

ENVIRONMENTAL HAZARDS: Do not apply directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands) and estuarine/marine habitats. Do not contaminate water by cleaning of equipment or disposal of wastes.

FOR CONTROL OF POWDERY MILDEW ON GREENHOUSE TOMATOES

NOTE TO BUYER/USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE

APPLICATIONS: The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than N. M. BARTLETT INC. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. N. M. BARTLETT INC. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used as described below.

Accordingly, the Buyer and User assume all liability arising, and agree to hold N. M. BARTLETT INC. harmless from any claims based on efficacy and/or phytotoxicity in connection with the use described below.

Wear goggles or a face shield, a NIOSH-approved respirator, chemical-resistant gloves, a hat, a long-sleeved shirt, long pants and rubber boots during mixing/loading, application, clean-up and repair activities. Do not enter or allow workers to enter into treated areas until 24 hours after application.

DIRECTIONS FOR USE

CROP/DISEASE

METHOD/RATE

GREENHOUSE TOMATO
Powdery Mildew

Apply 750 g per 1000 L water/ha; phi 1 day; maximum of 10 applications per crop cycle. Apply weekly from onset of first symptoms and during conditions favouring disease; two applications may be sufficient to control each incidence of disease; may cause slight foliar phytotoxicity.

FOR CONTROL OF POWDERY MILDEW ON GREENHOUSE PEPPERS

NOTE TO BUYER/USER : READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE

APPLICATIONS : The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than N.M. BARTLETT INC. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. N.M. BARTLETT INC. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used as described below.

Accordingly, the Buyer and User assume all liability arising, and agree to hold N.M. BARTLETT INC. harmless from any claims based on efficacy and/or phytotoxicity) in connection with the use described below.

Wear goggles or a face shield, a NIOSH-approved respirator, chemical-resistant gloves, a hat, a long-sleeved shirt, long pants and rubber boots during mixing/loading, application, clean-up and repair activities. Do not enter or allow workers to enter into treated areas until 24 hours after application.

DIRECTIONS FOR USE

CROP/DISEASE

METHOD/RATE

GREENHOUSE PEPPERS
Powdery Mildew

Apply 543-760g per 1000 L water/ha; apply as required; do not exceed 10 applications per crop cycle; apply on a minimum of 14 day interval.

**FOR CONTROL OF POWDERY MILDEW ON CANNABIS (MARIHUANA) PRODUCED
COMMERCIALY INDOORS**

NOTE TO BUYER/USER : READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE

APPLICATIONS : The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than N.M. BARTLETT INC. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. N.M. BARTLETT INC. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytoxicity) claims for this product when used as described below.

Accordingly, the Buyer and User assume all liability arising, and agree to hold N.M. BARTLETT INC. harmless from any claims based on efficacy and/or phytotoxicity) in connection with the use described below.

Wear goggles or a face shield, a NIOSH-approved respirator, chemical-resistant gloves, a hat, a long-sleeved shirt, long pants and rubber boots during mixing/loading, application, clean-up and repair activities. Do not enter or allow workers to enter into treated areas until 24 hours after application.

DIRECTIONS FOR USE

CROP/DISEASE

CANNABIS (marihuana)
produced commercially indoors
Powdery Mildew (*Sphaerotheca macularis*)

METHOD/RATE

Apply 750 g per 1000 L water/ha;; maximum of 10 applications per crop cycle. Apply weekly from onset of first symptoms and during conditions favouring disease; two applications may be sufficient to control each incidence of disease.

**N.M. BARTLETT INC., 4509 BARTLETT RD., BEAMSVILLE, ONTARIO L0R 1B1
1-905-563-8261 1-800-263-1287**